




e-ISSN: 2456-6632

This content is available online at AESA

Archives of Agriculture and Environmental Science

Journal homepage: www.aesacademy.org



ORIGINAL RESEARCH ARTICLE

Quantitative evaluation of essential oils for the identification of chemical constituents by gas chromatography/mass spectrometry

Ashish Uniyal^{a*}, Sachin N. Tikar^a, Om P. Agrawal^c, Devanathan Sukumaran^a and Vijay Veer^b

^aVector Management Division, Defence R & D Establishment, Gwalior-474002 (Madhya Pradesh), INDIA

^bDefence Research Laboratory, Tezpur - 784001 (Assam), INDIA

^cSchool of Studies in Zoology, Jiwaji University, Gwalior-474 011 (Madhya Pradesh), INDIA

*Corresponding author. E-mail: ashish.uniyal5@gmail.com

ARTICLE HISTORY

Received: 18 Sept. 2016

Revised received: 22 Sept. 2016

Accepted: 25 Sept. 2016

Keywords

Chemical constituents

Essential oils

Gas chromatography

Mass Spectrometry

ABSTRACT

Essential oils are greatly strenuous aromatic materials having various constituents. They are used in the preparation of various precious substances like making perfumes, medicines, cleaning agent, and aromatic treatment etc. The purpose of the present investigation was to identify the major and minor chemical constituent in eighteen essential oils *viz.*, amyris, basil, black pepper, camphor, catnip, chamomile, cinnamon, citronella, dill, frankincense, galbanum, jasmine, juniper, lavender, peppermint, rosemary, tagetes and thyme with the help of gas chromatography /mass spectrometry (GC/MS). In eighteen essential oils the identified compounds studied by GC-MS contain various types of high and low molecular weights of chemical ingredients. Therefore, GC/MS efficiently and speedily screened all the volatile elements present in the essential oils for the quantitative use of these identified chemical constituents for various reasons.

©2016 Agriculture and Environmental Science Academy

Citation of this article: Uniyal, A., Tikar, S.N., Agrawal, O.P., Sukumaran, D. and Veer, V. (2016). Quantitative evaluation of essential oils for the identification of chemical constituents by gas chromatography/mass spectrometry. *Archives of Agriculture and Environmental Science*, 1(1): 22-36.

INTRODUCTION

Essential oils are natural secondary plant products contain highly volatile liquid contain many chemicals. Most of the essential oils grouped into major chemical categories like alkaloids, phenolics and terpenoids are major constituents of essential oils. Essential oils are extracted from various parts of plants with the help of distillation methods such as hydro distillation, steam distillation hydro-steam distillation and solvent extraction. A variety of biological activities including behavioural responses have been recorded towards such essential oils in different living organisms. The response of biological agents comprises all activities that these volatile compounds or their mixtures exert on humans, animals, and other plants (Marcelle Gonny *et al.*, 2006; Suleiman Afsharypuor and Nahid Azarbajeny, 2006; Baser and Buchbauer, 2010).

The important oil producing plants are represented in more than thirty families of plants, comprising some ninety species. The majority of spices (cardamom, clove, nutmeg, ginger etc.) originate from tropical countries. Conversely the majority of herbs (bay, cumin, dill, marjoram, fennel, lavender, rosemary, thyme, etc.) grow in temperate

climates. The same plant grown in different regions and under different conditions can produce essential oils of widely diverse characteristics, which are known as "chemotypes". Common thyme (*Thymus vulgaris*) for example produces several chemotypes depending on the conditions of its growth and dominant constituent, notably the citral or linalool types, and the thymol or carvacrol type. It is therefore important not only to know the botanical name of the plant from which an oil has been produced, but also its place of origin and main constituents which are concerned to define its qualities (Lawless, 1992).

Essential oils are extracted from almost every conceivable plant part, such as flowers like rose and chamomile, leaves as peppermint and rosemary, fruits of orange and lemon, seeds as in coriander and fennel, grasses like lemongrass and ginger grass, roots and rhizomes as ginger, wood of cedar wood and sandalwood, bark like in cinnamon, gum as in frankincense (Tisserand, 1990). There are also essential oil from bulbs like garlic, dried flower buds like clove, and from stems or twig like clove stem. Usually they are liquid but can also be solid or semisolid, according to temperature such as rose. The majority of essential oils are clear or pale yellow in color, although a few are deeply

colored like German chamomile (blue). They are damaged by the effects of light, heat, air and moisture, and should always be kept in a cool environment, in tightly Stoppard dark glass bottles. Essential oils are dissolved in pure alcohol, fats and are not soluble in water (Tisserand, 1990; Suleiman Afsharypuor and Nahid Azarbayejany, 2006). The chemical constituents present in the different essential oils are used in perfumes, cosmetics, soaps, for flavoring foods, drinks and household cleaning products and various essential oils have been used in medical formulations. Keeping above in view, the present investigation was carried out this study, to determine the various chemical constituents present in the different essential oils using gas chromatography /mass spectrometry (GC/MS) and identified with their retention time and molecular weight.

MATERIALS AND METHODS

Collection of essential oils: Eighteen essential oils namely amyris (*Amyris balsamifera*), basil (*Ocimum basilicum*), black pepper (*Piper nigrum*), camphor (*Cinnamomum camphora*), catnip (*Nepeta cataria*), chamomile (*Anthemis nobilis*), cinnamon (*Cinnamomum zeylanicum*), citronella (*Cymbopogon winterianus*), dill (*Anethum graveolens*), frankincense (*Boswellia carteri*), galbanum (*Ferula galbaniflua*), jasmine (*Jasminum grandiflorum*), juniper (*Juniperus communis*), lavender (*Lavendula angustifolia*), peppermint (*Mentha piperita*), rosemary (*Rosmarinus officinalis*), tagetes (*Tagetes minuta*), thyme (*Thymus serpyllum*) were obtained from the authentic source of trade/ Fragrance and Flavour Development Center (FFDC), Kannuj (Uttar Pradesh) for the identification of essential oils and chemical constituents.

Gas chromatography /Mass spectrometry analysis: Gas chromatography /mass spectrometry are standard equip-

ment used to analyze different chemical components present in essential oil. GC/MS is a method that combines the features of gas-liquid chromatography and mass spectrometry to identify different substances within a sample. The identification of component in 23 essential oils was analyzed by gas chromatography/ mass spectrometry (GC/MS). GC/MS analysis of the oil was carried out on an Agilent gas chromatograph (2880A) with a (30 m × 0.32mm × 0.25 μ ID) equipped with an Agilent mass selective detector in the electron impact mode (Ionization energy: 70 eV). Helium gas was used as the carrier gas at constant flow rate 2 ml/minute and an injection volume of 1 μl was employed (Split ratio of 100:1) injector temperature 250°C; ion-source temperature 260°C, capillary: 30 m × 230μm, film thickness 0.25 μm, average velocity 44.6 cm/s, pressure 1.9 psi, purge flow 3 ml/minute, purge time 0.20 minute. The oven temperature was programmed from 50 to 325°C (isothermal for 2 minutes) with an increase of 5°C/minute to 160°C, then 20°C/minute to 260°C, equilibration time 1 minute, ramp 5°C/min, ending with a 30 minutes isothermal at 290°C. Total GC running time was 30 minutes. The volume of injected specimen of 1μl of diluted oil in methanol solution (1%).

Identification of chemical constituents: The chemical constituents of essential oils were identified in comparison with their retention indices. Identification of components of essential oil was based on retention indices (RI) and computer matching with the PBM libraries.

RESULTS AND DISCUSSION

Essential oils are highly concentrated aromatic substances found in plants contain numbers of organic constituents including hormones, vitamin and other natural elements. Therefore, it is important to identification the components of essential oils using gas chromatography /mass spectrometry

Table 1. Major chemical components of different essential oils identified by GC/MS.







| Oil | Chemical Name | R.T. | IUPAC Name | Molecular Formula | M.W. | Qual. % | Structure |
|--------|---------------|--------|---|-----------------------------------|------|---------|---|
| | eremophilene | 20.917 | 4a,5-dimethyl-3-prop-1-en-2-yl-2,3,4,5,6,7-hexahydro-1H-naphthalene | C ₁₅ H ₂₄ | 204 | 90 |  |
| | β-maaliene | 22.593 | 1,1,3a,7-tetramethyl-1a,2,3,3a,4,5,6,7b-octahydro-1H-cyclopropa(a)naphthalene | C ₁₅ H ₂₄ | 204 | 95 |  |
| | β-amorphene | 22.666 | 2-Isopropyl-5-methyl-9-methylene-bicyclo-1-decene(4.4.0) | C ₁₅ H ₂₄ | 204 | 93 |  |
| Amyris | β-cadinene | 22.824 | 4-isopropyl-1,6-dimethyldecahydronaphthalene | C ₁₅ H ₂₄ | 204 | 97 |  |
| | (+)-calarene | 23.296 | 1,1,7,7a-tetramethyl-2,3,5,6,7,7b-hexahydro-1aH-cyclopropa(a)naphthalene | C ₁₅ H ₂₄ | 204 | 97 |  |
| | driminol | 24.928 | 2,5,5,8a-tetramethyl-1,4,4a,6,7,8-hexahydronaphthalen-1-yl) | C ₁₅ H ₂₆ O | 222 | 94 |  |

Table 1. Contd.

| | | | | | | | |
|---------------------|------------------------------|--------|--|--|-----|----|--|
| | linalool oxide | 8.483 | 2-Methyl-2-vinyl-5-(1-hydroxy-1-methylethyl)tetrahydrofuran | C ₁₀ H ₁₈ O ₂ | 170 | 80 | |
| | linalool oxide | 8.926 | 2-Methyl-2-vinyl-5-(1-hydroxy-1-methylethyl)tetrahydrofuran | C ₁₀ H ₁₈ O ₂ | 170 | 90 | |
| Basil | linalool | 9.277 | 3, 7-dimethylocta-1, 6-dien-3-ol | C ₁₀ H ₁₈ O | 154 | 93 | |
| | estragole | 11.949 | 1-allyl-4-methoxybenzene | C ₁₀ H ₁₂ O | 148 | 98 | |
| | α -thujene | 4.854 | 4-methyl-1-propan-2-ylbicyclo(3.1.0)hex-3-ene | C ₁₀ H ₁₆ | 136 | 93 | |
| | β -pinene | 5.007 | 6,6-Dimethyl-2-methylenebicyclo(3.1.1)heptanes | C ₁₀ H ₁₆ | 136 | 97 | |
| | sabinene | 5.91 | 4-methylene-1-(1-methylethyl)bicyclo(3.1.0)hexane | C ₁₀ H ₁₆ | 136 | 97 | |
| | β -pinene | 6.019 | 6,6-Dimethyl-2-methylenebicyclo(3.1.1)heptanes | C ₁₀ H ₁₆ | 136 | 97 | |
| Black Pepper | δ -3-carene | 6.781 | 3,7,7-trimethylbicyclo(4.1.0)hept-3-ene | C ₁₀ H ₁₆ | 136 | 97 | |
| | DL-limonene | 7.371 | 1-Methyl-4-(1-methylethenyl)-cyclohexene | C ₁₀ H ₁₆ | 136 | 99 | |
| | <i>P</i> -mentha 1,4(8)-dine | 15.598 | 1-methyl-4-propan-2-ylidenecyclohexene | C ₁₀ H ₁₆ | 136 | 91 | |
| | (-)- α -copaene | 16.663 | 8-isopropyl-1,3-dimethyltricyclo(4.4.0.0 ^{2,7})dec-3-ene | C ₁₅ H ₂₄ | 204 | 99 | |
| | β -elemene | 17.005 | 2,4-Disopropenyl-1-methyl-1-Venylcyclohexane | C ₁₅ H ₂₄ | 204 | 99 | |
| | β -caryophyllene | 17.733 | 4, 11, 11-trimethyl-8-methylene-bicyclo(7.2.0)undec-4-ene | C ₁₅ H ₂₄ | 204 | 99 | |

Table 1. *Contd.*

| | | | | | | | |
|---------------------|-----------------------|--------|--|-----------------|-----|----|--|
| | α -bergamotene | 18.103 | 2,6-dimethyl-6-(4-methylpent-3-en-1-yl)bicyclo(3.1.1)hept-2-ene | $C_{15}H_{24}$ | 204 | 91 | |
| | α -humulene | 18.63 | 2,6,6,9-Tetramethyl-1,4,8-cycloundecatriene | $C_{15}H_{24}$ | 204 | 97 | |
| | curcumene | 19.308 | 1-Methyl-4-(6-methyl-5-hepten-2-yl)benzene | $C_{15}H_{22}$ | 204 | 96 | |
| | β -selinene | 19.447 | Eudesma-4(14),11-diene | $C_{15}H_{24}$ | 204 | 99 | |
| Black Pepper | α -selinene | 19.623 | Eudesma-3,11-diene | $C_{15}H_{24}$ | 204 | 97 | |
| | β -bisabolene | 19.658 | 1-Methyl-4-(6-methylhepta-1,5-dien-2-yl)cyclohex-1-ene | $C_{15}H_{24}$ | 204 | 97 | |
| | eremophilene | 20.942 | 4,5-dimethyl-3-prop-1-en-2-yl-2,3,4,5,6,7-hexahydro-1H-naphthalene | $C_{15}H_{24}$ | 204 | 90 | |
| | caryophyllene oxide | 21.676 | 4,12,12-trimethyl-9-methylene-5-oxatricyclo(8.2.0.04,6)dodecane | $C_{15}H_{24}O$ | 220 | 94 | |
| | ethylbenzene | 3.16 | Ethylbenzene | C_8H_{10} | 106 | 94 | |
| | m-xylol | 3.765 | 1,3-Dimethylbenzol | C_8H_{10} | 106 | 97 | |
| | O-xylene | 4.17 | 1,2-Dimethylbenzol | C_8H_{10} | 106 | 97 | |
| Camphor | camphene | 5.366 | 2,2-dimethyl-3-methylene-bicyclo(2.2.1)heptanes | $C_{10}H_{16}$ | 136 | 97 | |
| | δ -3-carane | 5.549 | 3,7,7-trimethylbicyclo(4.1.0)hept-3-ene | $C_{10}H_{16}$ | 138 | 58 | |
| | isocamphane | 5.865 | 2,2,3-Trimethylbicyclo(2.2.1)heptanes | $C_{10}H_{18}$ | 138 | 97 | |
| | norbornane | 6.045 | Bicyclo(2.2.1)heptanes | C_7H_{12} | 138 | 96 | |

Table 1. Contd.

| | | | | | | | |
|----------------|------------------------------|--------|--|--|-----|----|--|
| | <i>p</i> -menthane-3,8-diol | 6.364 | 2-(1-Hydroxy-1-methylethyl)-5-methylcyclohexanol | C ₁₀ H ₂₀ O ₂ | 140 | 74 | |
| | <i>p</i> -cimene | 7.198 | 1-Methyl-4-(1-methylethyl) benzene | C ₁₀ H ₁₄ | 134 | 97 | |
| | fencone | 8.944 | 1,3,3-Trimethylbicyclo(2.2.1) heptan-2-one | C ₁₀ H ₁₆ O | 154 | 95 | |
| Camphor | acetaldehyde | 9.422 | Ethanal | C ₂ H ₄ O | 152 | 90 | |
| | camphor | 10.547 | 1,7,7-Trimethylbicyclo(2.2.1) heptan-2-one | C ₁₀ H ₁₆ O | 152 | 98 | |
| | 2-methyl-4-nitrosoresorcinol | 15.828 | 2-Methyl-4-nitrosoresorcinol | C ₇ H ₇ NO ₃ | 153 | 50 | |
| | dimethyl phalate | 18.508 | dimethyl benzene-1,2-dicarboxylate | C ₁₀ H ₁₀ O ₄ | 194 | 94 | |
| | linalool | 9.268 | 3, 7-dimethylocta-1, 6-dien-3-ol | C ₁₀ H ₁₈ O | 154 | 91 | |
| | β -citronellol | 12.794 | 3,7-Dimethyloct-6-en-1-ol | C ₁₀ H ₂₀ O | 156 | 98 | |
| | geraniol | 13.434 | 3,7-Dimethyl-2,6-octadien-1-ol | C ₁₀ H ₁₈ O | 154 | 94 | |
| | neral | 14.012 | 3,7-dimethylocta-2,6-dienal | C ₁₀ H ₁₆ O | 152 | 49 | |
| Catnip | 2,6-Octadine,2,6-Dimethyl | 15.816 | 2,6-Octadine,2,6-Dimethyl | C ₁₀ H ₁₈ | 138 | 92 | |
| | 2,6-Octadine,2,6-Dimethyl | 16.058 | 2,6-Octadine,2,6-Dimethyl | C ₁₀ H ₁₈ | 138 | 98 | |
| | butanoic acid | 16.282 | Butanoic acid | C ₄ H ₈ O ₂ | 224 | 91 | |
| | 4-cyclopropyl cyclohexane | 16.702 | 4-Cyclopropyl Cyclohexane | C ₉ H ₁₆ | 112 | 58 | |

Table 1. *Contd.*

















| | | | | | | | |
|-------------------|------------------------------|--------|---|--|-----|----|---|
| | geraniol ester | 16.807 | 2,6-Octadien-1-ol, 3,7-dimethyl-, acetate | C ₁₀ H ₁₈ O | 196 | 91 |  |
| | caryophyllene | 17.718 | 4, 11, 11-trimethyl-8-methylene-bicyclo (7.2.0) undec-4-ene | C ₁₅ H ₂₄ | 204 | 99 |  |
| Catnip | 1-nitrocyclohexene | 19.459 | Nitrocyclohexane | C ₆ H ₁₁ NO ₂ | 127 | 50 |  |
| | β -caryophyllene oxide | 21.668 | 4,12,12-trimethyl-9-methylene-5-oxatricyclo(8.2.0.04,6)dodecane | C ₁₅ H ₂₄ O | 220 | 94 |  |
| | tricyclene | 4.768 | 1,4,7,10-tetraazacyclododecane | C ₁₀ H ₁₆ | 136 | 50 |  |
| | α -pinene | 5.003 | 2, 6, 6-Trimethyl bicycle (3.1.1) hept-2-ene | C ₁₀ H ₁₆ | 136 | 86 |  |
| | camphene | 5.366 | 2,2-dimethyl-3-methylene-bicyclo (2.2.1)heptanes | C ₁₀ H ₁₆ | 136 | 94 |  |
| | β -pinene | 6.016 | 6,6-Dimethyl-2-methylenebicyclo (3.1.1)heptanes | C ₁₀ H ₁₆ | 136 | 94 |  |
| Chamo-mile | cymene | 7.193 | 1-Methyl-4-(1-methylethyl) benzene | C ₁₀ H ₁₄ | 134 | 87 |  |
| | 1,8-cineole | 7.428 | 1,3,3-Trimethyl-2-oxabicyclo (2,2,2)octane | C ₁₀ H ₁₈ O | 154 | 98 |  |
| | δ -3-carene | 7.781 | 3,7,7-trimethylbicyclo(4.1.0)hept-3-ene | C ₁₀ H ₁₆ | 136 | 50 |  |
| | neryl acetate | 9.274 | 3,7-Dimethyl-2,6-octadien-1-yl acetate | C ₁₂ H ₂₀ O ₂ | 196 | 53 |  |
| | α -terpinene | 11.882 | 4-Methyl-1-(1-methylethyl)-1,3-cyclohexadiene | C ₁₀ H ₁₆ | 136 | 60 |  |
| | β -farnesene | 18.633 | 3,7,11-trimethyldodeca-1,3,6,10-tetraene | C ₁₅ H ₂₄ | 204 | 96 |  |
| | 1,6-cyclodecadiene | 19.252 | 1,6-Cyclodecadiene | C ₁₀ H ₁₆ | 204 | 94 |  |
| | Cis α -bisabolene | 19.784 | 4-[(1Z)-1,5-Dimethyl-1,4-hexadienyl]-1-methyl-1-cyclohexene | C ₁₅ H ₂₄ | 204 | 90 |  |

Table 1. Contd.

| | | | | | | | |
|-------------------|---------------------------------|--------|--|-------------------|-----|----|--|
| Chamomile | α -franesene | 19.892 | 3,7,11-trimethyldodeca-1,3,6,10-tetraene | $C_{15}H_{24}$ | 204 | 91 | |
| | <i>Trans</i> -caryophyllene | 19.948 | 4,11,11-trimethyl-8-methylene-bicyclo(7.2.0)undec-4-ene | $C_{15}H_{24}$ | 204 | 72 | |
| | α -longipinene | 20.01 | 2,6,6,9-Tetramethyltricyclo(5.4.0.0 ^{2,8})undec-9-ene | $C_{15}H_{24}$ | 204 | 64 | |
| | <i>Cis</i> α -bisabolene | 20.73 | 4-[(1 <i>Z</i>)-1,5-Dimethyl-1,4-hexadienyl]-1-methyl-cyclohexene | $C_{15}H_{24}$ | 204 | 93 | |
| | cinnamal | 13.966 | 2-Propenal, 3-phenyl | $C_{15}H_{20}O$ | 132 | 98 | |
| Cinnamon | β -isosafrole | 14.403 | 5-[(1 <i>E</i>)-Prop-1-en-1-yl]-1,3-benz | $C_{10}H_{10}O_2$ | 162 | 49 | |
| | <i>p</i> -eugenol | 16.054 | 2-methoxy-4-prop-2-enylphenol | $C_{10}H_{12}O_2$ | 164 | 98 | |
| | β -caryophyllene | 17.721 | 4, 11, 11-trimethyl-8-methylene-bicyclo (7.2.0) undec-4-ene | $C_{15}H_{24}$ | 204 | 90 | |
| | cinnamaldehyde | 18.191 | (2 <i>E</i>)-3-phenylprop-2-enal | C_9H_8O | 230 | 93 | |
| | 2-propen-1-ol | 18.446 | 2-Propen-1-ol | C_3H_6O | 176 | 53 | |
| Citronella | α -humulene | 18.621 | 2,6,6,9-Tetramethyl-1,4-8-cycloundecatriene | $C_{15}H_{24}$ | 204 | 39 | |
| | DL-limonene | 7.306 | 1-Methyl-4-(1-methylethenyl)-cyclohexene | $C_{10}H_{16}$ | 136 | 99 | |
| | α -terpinolene | 9.297 | 1-Methyl-4-(propan-2-ylidene)cyclohex-1-ene | $C_{10}H_{16}$ | 136 | 91 | |
| | isopulegol | 10.602 | 5-methyl-2-prop-1-en-2-ylcyclohexan-1-ol | $C_{10}H_{18}O$ | 154 | 98 | |
| | citronella | 10.712 | 3,7-dimethyloct-6-enal | $C_{10}H_{18}O$ | 154 | 98 | |

Table 1. Contd.

| | | | | | | | |
|-------------------|------------------------------|--------|--|-----------------------------------|-----|----|--|
| | camphene | 10.859 | 2,2-dimethyl-3-methylene-bicyclo(2.2.1)heptanes | C ₁₀ H ₁₆ | 136 | 91 | |
| | β -citronellol | 12.799 | 3,7-Dimethyloct-6-en-1-ol | C ₁₀ H ₂₀ O | 156 | 98 | |
| | geraniol | 13.44 | 3,7-Dimethyl-2,6-octadien-1-ol | C ₁₀ H ₁₈ O | 154 | 87 | |
| Citronella | farnesol L | 15.631 | 3,7,11-trimethyldodeca-2,6,10-trien-1-ol | C ₁₅ H ₂₆ O | 222 | 49 | |
| | geranyl propionate | 16.803 | 2,6-Octadien-1-ol, 3,7-dimethyl-, propanoate | C ₁₅ H ₂₄ | 210 | 90 | |
| | β -elemene | 17.001 | 2,4-Disopropenyl-1-methyl-1-Venylcyclohexane | C ₁₅ H ₂₄ | 204 | 95 | |
| | δ -cadinene | 20.175 | 4,7-dimethyl-1-(propan-2-yl)-1,2,3,5,6,8a-hexahydronaphthalene | C ₁₅ H ₂₄ | 204 | 99 | |
| | (+)-longifolene | 20.928 | 3,3,7-trimethyl- 8-methylenetricyclo- (5.4.0.0) undecane | C ₁₅ H ₂₄ | 204 | 95 | |
| | α -pinene | 4.981 | 2, 6, 6-Trimethyl bicycle (3.1.1) hept-2-ene | C ₁₀ H ₁₆ | 136 | 95 | |
| | β -myrcene | 5.856 | 7-Methyl-3-methylene-1,6-octadiene | C ₁₀ H ₁₆ | 134 | 94 | |
| | <i>O</i> -cymol | 7.177 | 1-methyl-2-propan-2-ylbenzene | C ₁₀ H ₁₄ | 134 | 97 | |
| | DL-limonene | 7.298 | 1-Methyl-4-(1-methylethenyl)-cyclohexene | C ₁₀ H ₁₆ | 136 | 99 | |
| Dill | 2-Methyl-1-propenyl, benzene | 8.928 | 2-methylprop-1-enylbenzene | C ₁₀ H ₁₂ | 132 | 97 | |
| | limonene epoxide | 10.264 | 6-methyl-3-prop-1-en-2-yl-7-oxabicyclo(4.1.0) heptanes | C ₁₀ H ₁₆ O | 152 | 76 | |
| | limonene oxide | 10.527 | 6-methyl-3-prop-1-en-2-yl-7-oxabicyclo(4.1.0) heptanes | C ₁₀ H ₁₆ O | 134 | 78 | |
| | (+)-dihydrocarvone | 11.95 | (2R,5R)-5-Isopropenyl-2-methylcyclohexanone | C ₁₀ H ₁₆ O | 152 | 99 | |
| | cyclohexanone | 12.124 | Cyclohexanone | C ₆ H ₁₀ O | 152 | 98 | |

Table 1. Contd.

| | | | | | | | |
|---------------------|------------------------|---|--|--|-----|----|--|
| Dill | carveol | 12.524 | 2-Methyl-5-(1-methylethenyl)-2-cyclohexen-1-ol | C ₁₀ H ₁₆ O | 152 | 55 | |
| | (+)-carvone | 13.211 | 5-Isopropenyl-2-methyl-2-cyclohexenone, <i>p</i> -Mentha-6,8-dien-2-one | C ₁₀ H ₁₄ O | 150 | 97 | |
| | thymol | 14.489 | 2-Isopropyl-5-methylphenol | C ₁₀ H ₁₄ O | 150 | 95 | |
| | dillapiole | 22.536 | 1-Allyl-2,3-dimethoxy-4,5-(methylenedioxy)benzene | C ₁₂ H ₁₄ O ₄ | 222 | 97 | |
| | α -thujene | 4.85 | 4-Methyl-1-(propan-2-yl)bicycle(3.1.0)hexan-3-one | C ₁₀ H ₁₆ O | 134 | 93 | |
| | α -pinene | 5.006 | 2, 6, 6-Trimethyl bicycle (3.1.1) hept-2-ene | C ₁₀ H ₁₆ | 136 | 97 | |
| | sabinene | 5.907 | 4-methylene-1-(1-methylethyl) bicycle(3.1.0) hexane | C ₁₀ H ₁₆ | 136 | 96 | |
| | β -pinene | 6.06 | 6,6-Dimethyl-2-methylenebicyclo(3.1.1) heptanes | C ₁₀ H ₁₆ | 136 | 96 | |
| β -ocimene | 6.312 | 3,7-dimethylocta-1,3,6-triene | C ₁₀ H ₁₆ | 136 | 45 | | |
| γ -terpinene | 6.778 | 1-Isopropyl-4-methyl-1,4-cyclohexadiene | C ₁₀ H ₁₆ | 136 | 87 | | |
| <i>p</i> -cimene | 7.197 | 1-Methyl-4-(1-methylethyl) benzene | C ₁₀ H ₁₄ | 134 | 90 | | |
| Frankincense | DL-limonene | 7.306 | 1-Methyl-4-(1-methylethenyl)-cyclohexene | C ₁₀ H ₁₆ | 136 | 93 | |
| | thujopsene | 8.765 | 2,4a,8,8-Tetramethyl-1,1a,4,4a,5,6,7,8-octahydrocyclopropano(d)naphthalene | C ₁₅ H ₂₄ | 204 | 38 | |
| | α -thujone | 9.74 | Methyl-1-(propan-2-yl)bicycle(3.1.0)hexan-3-one | C ₁₀ H ₁₆ O | 134 | 90 | |
| | <i>p</i> -allylanisole | 11.934 | 1-allyl-4-methoxybenzene | C ₁₀ H ₁₂ O | 148 | 96 | |

Table 1. Contd.















| | | | | | | | |
|-----------------|-----------------------|--------|---|--|-----|----|---|
| | tricyclene | 4.755 | 6,7,7-trimethyl-2,3,4,5-tetrahydro-1H-tricyclo(2.2.1.0 ^{2,6})heptanes | C ₁₀ H ₁₆ | 136 | 96 |  |
| | α -pinene | 4.991 | 2,6,6-Trimethyl bicyclo(3.1.1)hept-2-ene | C ₁₀ H ₁₆ | 136 | 97 |  |
| | camphene | 5.357 | 2,2-dimethyl-3-methylenebicyclo(2.2.1)heptanes | C ₁₀ H ₁₆ | 136 | 98 |  |
| | β -cimene | 5.856 | 1-methyl-3-propan-2-ylbenzene | C ₁₀ H ₁₄ | 134 | 94 |  |
| | β -pinene | 6.005 | 6,6-Dimethyl-2-methylenebicyclo(3.1.1)heptane | C ₁₀ H ₁₆ | 136 | 97 |  |
| Galbanum | δ -3-carene | 6.765 | 3,7,7-trimethylbicyclo(4.1.0)hept-3-ene | C ₁₀ H ₁₆ | 136 | 97 |  |
| | <i>o</i> -cimene | 7.182 | 1-Menthyl-2-isopropylbenzene | C ₁₀ H ₁₄ | 134 | 97 |  |
| | DL-limonene | 7.298 | 1-Methyl-4-(1-methylethenyl)cyclohexene | C ₁₀ H ₁₆ | 136 | 99 |  |
| | pinocarveol | 10.355 | 6,6-Dimethyl-2-methylenebicyclo(3.1.1)heptan-3-ol | C ₁₀ H ₁₆ O | 152 | 56 |  |
| | mentha-1,4,8-triene | 10.527 | 1-methyl-4-prop-1-en-2-ylcyclohexa-1 | C ₁₀ H ₁₄ | 134 | 78 |  |
| | (-) myrtenol | 11.895 | 6,6-Dimethylbicyclo(3.1.1)hept-2-ene-2-methanol | C ₁₀ H ₁₆ O | 152 | 97 |  |
| Jasmine | α -toluenol | 7.496 | Phenylmethenol | C ₇ H ₈ O | 108 | 97 |  |
| | α -terpinolene | 9.227 | 1-Methyl-4-(propan-2-ylidene)cyclohex-1-ene | C ₁₀ H ₁₆ | 136 | 90 |  |
| | eugenol | 16.5 | 4-Allyl-2-methoxyphenol | C ₁₀ H ₁₂ O ₂ | 164 | 98 |  |

Table 1. Contd.


















| | | | | | | | |
|-----------------|---------------------|---------|--|--|-----|----|---|
| | ascabin | 24.899 | Benzyl benzoate | C ₁₄ H ₁₂ O ₂ | 212 | 98 |  |
| Jasmine | palmitic acid | 26.2772 | hexadecanoic acid | C ₁₆ H ₃₂ O ₂ | 270 | 96 |  |
| | isophytol | 26.417 | 3,7,11,15-Tetramethyl-1-hexadecen-3-ol | C ₂₀ H ₄₀ O | 296 | 86 |  |
| | phytol | 27.407 | 3,7,11,15-tetramethyl-2-hexadecen-1-ol | C ₂₀ H ₄₀ O | 290 | 43 |  |
| | neophytadiene | 27.937 | 7,11,15-trimethyl-3-methylidenehexadec-1-ene | C ₂₀ H ₃₈ | 278 | 90 |  |
| | <i>p</i> -ocimene | 4.838 | 3,7-Dimethyl-1,3,6-Octatriene | C ₁₀ H ₁₆ | 138 | 83 |  |
| | α -pinene | 5.00 | 2, 6, 6-Trimethyl bicycle (3.1.1) hept-2-ene | C ₁₀ H ₁₆ | 136 | 97 |  |
| Juniper | sabinene | 5.898 | 4-methylene-1-(1-methylethyl) bicycle(3.1.0)hexane | C ₁₀ H ₁₆ | 136 | 91 |  |
| | β -pinene | 6.011 | 6,6-Dimethyl-2-methylenebicyclo(3.1.1) heptanes | C ₁₀ H ₁₆ | 136 | 97 |  |
| | γ -terpinene | 6.769 | 1-Isopropyl-4-methyl-1,4-cyclohexadiene | C ₁₀ H ₁₆ | 136 | 90 |  |
| | DL-limonene | 7.309 | 1-Methyl-4-(1-methylethenyl)-cyclohexene | C ₁₀ H ₁₆ | 136 | 98 |  |
| | β -Cymene | 7.188 | 1-1-methyl-3-propan-2-ylbenzene | C ₁₀ H ₁₄ | 134 | 87 |  |
| | limonene oxide | 9.199 | 6-methyl-3-prop-1-en-2-yl-7-oxabicyclo(4.1.0)heptane (Click) | C ₁₀ H ₁₆ O | 152 | 53 |  |
| Lavender | para- mentha-triene | 9.505 | 1-methyl-4-prop-1-en-2-ylcyclohexa-1,3-diene | C ₁₀ H ₁₄ | 134 | 22 |  |
| | camphor | 10.536 | 1,7,7-Trimethylbicyclo(2.2.1) heptan-2-one | C ₁₀ H ₁₆ O | 152 | 72 |  |
| | benzyl acetate | 10.978 | Benzyl acetate | C ₉ H ₁₀ O ₂ | 150 | 64 |  |
| | δ -3-carene | 13.395 | 3,7,7-trimethylbicyclo(4.1.0) hept-3-ene | C ₁₀ H ₁₆ | 136 | 94 |  |

Table 1. Contd.

| | | | | | | | |
|-------------------|---------------|--------|--|-----------------------------------|-----|----|--|
| | camphene | 10.597 | 2,2-dimethyl-3-methylene-bicyclo(2.2.1)heptanes | C ₁₀ H ₁₆ | 136 | 93 | |
| | cyclohexanone | 10.794 | Cyclohexanone | C ₆ H ₁₀ O | 154 | 98 | |
| | DL-menthol | 11.182 | 2-Isopropyl-5-methylcyclohexanol | C ₁₀ H ₂₀ O | 156 | 91 | |
| Peppermint | L-(-) menthol | 11.419 | 2-Isopropyl-5-methylcyclohexanol, 5-Methyl-2-(1-methylethyl)cyclohexanol | C ₁₀ H ₂₀ O | 156 | 91 | |
| | 3-p-menthanol | 11.685 | 3-p-Menthanol | C ₁₀ H ₂₀ O | 156 | 91 | |
| | norcarane | 14.468 | Bicyclo(4,1,0)heptanes | C ₇ H ₁₂ | 136 | 96 | |
| | α-pinene | 5.004 | 2, 6, 6-Trimethyl bicycle (3.1.1) hept-2-ene | C ₁₀ H ₁₆ | 136 | 97 | |
| | camphene | 5.37 | 2,2-dimethyl-3-methylene-bicyclo(2.2.1)heptane | C ₁₀ H ₁₆ | 136 | 98 | |
| Rosemary | β-pinene | 6.011 | 6,6-Dimethyl-2-methylenebicyclo(3.1.1) heptanes | C ₁₀ H ₁₆ | 136 | 96 | |
| | cymol | 7.197 | 2,6-dimethyl-3-propan-2-ylpheno | C ₁₀ H ₁₄ | 134 | 95 | |
| | DL-limonene | 7.31 | 1-Methyl-4-(1-methylethenyl)-cyclohexene | C ₁₀ H ₁₆ | 136 | 99 | |
| | 1,8-cineole | 7.441 | 1,3,3-Trimethyl-2-oxabicyclo (2,2,2)octane | C ₁₀ H ₁₈ O | 154 | 99 | |
| | α-terpinolene | 9.274 | Methyl-4-(propan-2-ylidene) cyclohex-1-ene | C ₁₀ H ₁₆ | 136 | 78 | |
| | camphor | 10.55 | 1,7,7-Trimethylbicyclo(2.2.1) heptan-2-one | C ₁₀ H ₁₆ O | 152 | 98 | |

Table 1. Contd.





















| | | | | | | | |
|--------------------|------------------------------|--|---|--|-----|---|---|
| Rosemary | isobroneol | 10.993 | 4,7,7-trimethylbicyclo(2.2.1) heptan-3-ol | C ₁₀ H ₁₆ O | 154 | 86 |  |
| | α -pinene | 11.892 | 2, 6, 6-Trimethyl bicycle (3.1.1) hept-2-ene | C ₁₀ H ₁₆ | 136 | 87 |  |
| | pichtosine | 14.359 | Bicyclo(2.2.1)heptan-2-ol, 1,7,7-trimethyl-, acetate | C ₁₂ H ₂₀ O ₂ | 196 | 91 |  |
| | β -caryophyllene oxide | 21.667 | 4,12,12-trimethyl-9-methylene-5-oxatricyclo(8.2.0.0 ^{4,6}) dodecane | C ₁₅ H ₂₄ O | 220 | 87 |  |
| | α -pinene | 4.999 | 2, 6, 6-Trimethyl bicycle (3.1.1) hept-2-ene | C ₁₀ H ₁₆ | 136 | 95 |  |
| | β -pinene | 6.001 | 6,6-Dimethyl-2-methylenebicyclo(3.1.1) heptanes | C ₁₀ H ₁₆ | 136 | 94 |  |
| Tagetes | <i>p</i> -cymene | 7.188 | 1-Methyl-4-(1-methylethyl) benzene | C ₁₀ H ₁₄ | 134 | 86 |  |
| | DL-limonene | 7.311 | 1-Methyl-4-(1-methylethenyl)-cyclohexene | C ₁₀ H ₁₆ | 136 | 99 |  |
| | 1,8-cineole | 7.428 | 1,3,3-Trimethyl-2-oxabicyclo (2,2,2)octane | C ₁₀ H ₁₈ O | 154 | 52 |  |
| | dihydro-tagetone | 7.959 | 2,6-Dimethyloct-7-en-4-one | C ₁₀ H ₁₈ O | 154 | 78 |  |
| | linalool oxide trans | 8.483 | 2-Methyl-2-vinyl-5-(1-hydroxy-1-methylethyl)tetrahydrofuran | C ₁₀ H ₁₈ O ₂ | 170 | 72 |  |
| | furfuryl alcohol | 8.921 | 2-Furanmenthanol | C ₅ H ₆ O ₂ | 170 | 49 |  |
| δ -3-carene | 9.274 | 3,7,7-trimethylbicyclo(4.1.0) hept-3-ene | C ₁₀ H ₁₆ | 136 | 87 |  | |

Table 1. Contd.

| | | | | | | | |
|-------|------------------|--------|---|-----------------|-----|----|---|
| Thyme | α -pinene | 4.986 | 2, 6, 6-Trimethyl bicycle (3.1.1) hept-2-ene | $C_{10}H_{16}$ | 136 | 97 |  |
| | β -pinene | 5.997 | 6,6-Dimethyl-2-methylenebicyclo(3.1.1)heptanes | $C_{10}H_{16}$ | 136 | 97 |  |
| | cymene | 7.188 | 1-Methyl-4-(1-methylethyl) benzene | $C_{10}H_{14}$ | 134 | 97 |  |
| | DL-limonene | 7.297 | 1-Methyl-4-(1-methylethenyl)-cyclohexene | $C_{10}H_{16}$ | 136 | 97 |  |
| | moslene | 8.075 | 1,4-Cyclohexadiene, 1-methyl-4-(1-methylethyl) | $C_{10}H_{16}$ | 136 | 97 |  |
| | α -pinene | 11.092 | 2, 6, 6-Trimethyl bicycle (3.1.1) hept-2-ene | $C_{10}H_{16}$ | 136 | 42 |  |
| | carvacrol | 14.709 | 5-isopropyl-2-methylphenol 2-Methyl-5-(1-methylethyl)-phenol | $C_{10}H_{14}O$ | 150 | 94 |  |

try (GC/MS) for detection of major and trace constituents. In the present study, the identification of highly volatile chemical components present in essential oils with the help of GC/MS showed many monoterpenes such as hydrocarbons (α -Pinene), alcohol (geraniol, menthol, linalool), ethers (1,8-cineol), aldehydes (cinnamaldehyde) etc. The composition of essential oils is highly diverse in different plant species and chemical components present in essential oils and some chemical constituents are common in few essential oils. The chemical compositions of essential oil are different in different stages of plant development and the chemical components of essential oils especially monoterpenes directly depend on temperature, weather condition and soil acidity as earlier reported by Clark and Manery (1981).

The components of essential oils were analysed and identified by GC-MS using DB-5 fused silica capillary column (Table 1). The major constituent α -Pinene (97%) of Juniper was dominant monoterpene (Gonny *et al.*, 2005). A monocyclic monoterpene DL-Limonene 98 (R.T-7.311) is a major constituent of tagetes oil (Chamorro *et al.*, 2008) and Linalool 93% (RT- 9.277) is an acyclic monoterpene alcohol also known as 3, 7-dimethylocta-1, 6-dien-3-ol (Benedec *et al.*, 2009). Cinnamon essential oil contains Cinnamaldehyde 93% (RT- 18.191) as a major component and it is a monocyclic monoterpene alcohol

and also known as Cinnamic aldehyde or trans Cinnamaldehyde (98%) (Adams, 1989) and the dominant chemical constituent of chamomile oil is 1, 8-Cineole 98% (RT-7.428). 1, 8-Cineole or Eucalyptol is a bicyclic monoterpene alcohol also called 1, 8-Epoxy-*p*-menthane (Adams, 1989). In frankincense oil α -Pinene 97% (RT- 5.006) a bicyclic monoterpene is the dominant constituent. It is an alkene and it contains a reactive four-membered ring (Woolley *et al.*, 2012). Moreover, Linalool (96%) is the main component of lavender oil (Afsharypur and Azarbajany, 2006) and the major component in amyris essential oil is β -Cadinene 97% (RT- 22.824) also called Cadina-3,9-diene.

Chemically, the Cadinenes are bicyclic sesquiterpenes (Lawrence, 1990). However, from the dominant constituent of dill oil is (+)-Carvone 97% (RT- 13.211) (Radulescu *et al.*, 2010) also called *p*-Mentha-6,8-dien-2-one a monocyclic monoterpene ketone and DL-Limonene 99% (7.298) is the dominant constituents in galbanum oil a monocyclic monoterpene hydrocarbon (Adams, 1995). The major dominant component identified in catnip oil is Caryophyllene 99% (RT- 17.718). Caryophyllene or β Caryophyllene is a natural bicyclic sesquiterpene (Wesolowska *et al.*, 2011) and Thymol 95% (RT- 14.514) is the dominant constituent of thyme oil. Thymol is a monocyclic monoterpene alcohol (Ahmad *et al.*, 2006). The major

component of black pepper oil is β - Caryophyllene (99%) (Jirovetza *et al.*, 2012) and the major component of citronella oil is citronella (98%), citronella oil contains two derivatives such as alcohol citronellol and the aldehyde citronellal (Cassel and Vargas, 2006). Rosemary essential oil contains Camphene 98% (RT- 5.370) as a dominant constituent identified by GC/MS is a bicyclic monoterpene (Martinez *et al.*, 2009) and jasmine contain Eugenol (98%) (Adams, 1989). Moreover, menthol (91%) is the dominant chemical constituent in peppermint also known as 3-*p*-menthanol is a monocyclic monoterpene alcohol (Derwich *et al.*, 2010) and camphor oil contains camphor 98% (RT- 10.547) is abicyclic monoterpene as a main constituent (Guenther, 1950).

Therefore, the essential oils were analyzed using GC/MS to get some major constituents with their retention time and molecular weight with quality percentage but some constituents are common in some essential oils with slightly difference in retention time such as Linalool present in lavender and basil as dominant component with retention time ranging from 9.271RT to 9.277RT.

Conclusions

The present study concluded that several major chemical constituents were identified from different essential oils like amyris (*A. balsamifera*), basil (*O. basilicum*), black pepper (*P. nigrum*), camphor (*C. camphora*), catnip (*N. cataria*), chamomile (*A. nobilis*), cinnamon (*C. zeylanicum*), citronella (*C. winterianus*), dill (*A. graveolens*), frankincense (*B. carteri*), galbanum (*F. galbaniflua*), jasmine (*J. grandiflorum*), juniper (*J. communis*), lavender (*L. angustifolia*), peppermint (*M. piperita*), rosemary (*R. officinalis*), tagetes (*T. minuta*), thyme (*T. serpyllum*) using GC/MS. These essential oils have several chemical constituents and can be used for multiple purposes such as use in national and international market of flavor and fragrances, cosmetic industries, agricultural industries, household insecticides, pharmacy, alternative medicine, integrated pest management, etc. Therefore, GC/MS can be effectively used for the identification of different chemical constituents present in the various essential oils.

ACKNOWLEDGEMENTS

The authors are grateful to Prof. (Dr.) M.P. Kaushik, former Director, Defence Research & Development Establishment (D.R.D.E), Gwalior and all the colleagues in the Department of Vector Management Division for their help and support while carrying out this investigation on essential oils. Fragrance and Flavor Development Center (FFDC), Kannuj, Uttar Pradesh, India is also kindly acknowledged by the authors for providing essential oils.

Open Access: This is open access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author(s) and the source are credited.

REFERENCES

- Adams, R. P. (1989). Identification of essential oils by ion trap mass spectroscopy. Academic Press, New York, NY, USA.
- Adams, R. P. (1995). Identification of essential oils components by gas chromatography/mass spectroscopy. Allured Publ. Corp., Illinois.
- Ahmad, M. A., Khokhar, I., Ahmad, I., Kashmiri, A. M., Adnan, A. and Ahmad, M. (2006). Study of antimicrobial activity and composition by GC/MS spectroscopic analysis of the essential oil of *Thymus sperphyllum*. *Internet Journal of Food Safety*, 5: 56-60.
- Baser, C. H. K. and Buchbauer, G. (2010). Hand book of essential oils: Science, technology and applications. Raton Florida: CRC Press, Boca Raton, New York.
- Benedec, D., Oniga, I., Oprean, R. and Tames, M. (2009). Chemical composition of the essential oils of *ocimum basilicum* L. cultivated in Romania. *Farmica*, 57(5): 625-629.
- Clark, R. J. and Menary, R. C. (1981). Variations in composition of peppermint oil in relation to production areas. *Economic Botany*, 35: 59-69.
- Derwich, E., Benziane, Z., Taouil, R., Senhaji, O. and Touzani, M. (2010). Aromatic plants of Morocco: GC/MS analysis of the essential oils of leaves of *Mentha piperita*. *Advances in Environmental Biology*, 4(1): 80-85.
- Gonny, M., Cavaleiro, C., Salgueiro, L. and Casanova, J. (2005). Analysis of *Juniperus communis* subsp. *alpina* needle, berry, wood and root oils by combination of GC, GC/MS and 13C-NMR. *Flavour Fragrance Journal*, 21: 99-106.
- Guenther, E. (1950). 3RD Ed. *The Essential Oils*. D. Van Nostrand, New York.
- Jirovetza, L., Buchbauer, G., Ngassoumb, B. M. and Geissler, M. (2012). Aroma compound analysis of *Piper nigrum* and *Piper guineense* essential oils from Cameroon using solid-phase microextraction-gas chromatography, solid-phase micro extraction GC/MS and olfactometry. *Journal of Chromatography*, 976: 265-275.
- Lawrence, B. M. (1990). Amyris oil. *Progress in essential oils. Perfumer and Flavorist*, 15 (2): 78.
- Lawless, J. (1992). The encyclopedia of essential oils. Shaftesbury, Dorset. Rockport, Massachusetts.
- Marcelle Gonny, Carlos Cavaleiro, Ligia Salgueiro and Joseph Casanova (2006). Analysis of *Juniperus communis* subsp. *alpina* needle, berry, wood and root oils by combination of GC, GC/MS and 13C-NMR. *Flavour and Fragrance Journal*, 21: 99-106.
- Martinez, A. L., Gonzalez-Trujano, M.E., Pellicer, F., Lopez-Munoz, F.J. and Navarrete, A. (2009). Antinociceptive effect and GC/MS analysis of *Rosmarinus officinalis* L. essential oil from its aerial parts. *Planta Medicina*, 75: 508-511.
- Radulescu, V., Popescu, M. L. and Ilies, D. (2010). Chemical composition of the volatile oil from different plants parts of *Anthem graveolens* L. cultivated in Romania. *Farmica*, 58(5): 594-600.
- Suleiman Afsharypuor and Nahid Azarbajeny (2006). Chemical constituents of the flower essential oil of *Lavandula officinalis* Chaix. from Isfahan (Iran). *Iranian Journal of Pharmaceutical Sciences*, 2(3): 169-172.
- Tisserand, R. (1990). The essential oil safety data manual. Tisserand Aromatherapy Institute Brighton. Sussex. England.
- Wesolowska, A., Jadcak, D. and Gerzeszczuk, M. (2011). MS analysis of lemon catnip (*Nepeta cataria* L. var. *citriodora* Balbis) essential oil. *Acta Chromatographica*, 1: 169-180.
- Woolley, L.C., Suhailb, M. M., Smitha, B.L., Borena, E.K., Taylora, C.L., Schreudera, F.M., Chaia, K. J., Haq, S.C.H., Linf, K.H., Al-Shahrig, A.A., Saif, Al-Hatmih and Younga, G. D. (2012). Chemical differentiation of *Boswellia sacra* and *Boswellia carterii* essential oils by gas chromatography and chiral GC/MS. *Journal of Chromatography*, 1261: 158-163.